

Testing the quantum arrow of time in weak decays

A. J. Bevan, G. Inguglia, and M. Zoccali

Queen Mary, University of London, Mile End Road, E1 4NS, United Kingdom

(Dated: February 19, 2013)

The arrow of time is a familiar concept; entropy increases, landscapes are eroded away by the elements, and we inevitably grow older. These irreversible macroscopic effects taint our intuition of time, however it is well known that the quantum arrow of time is different from this entropic behaviour. The laws of quantum physics can be studied under the mathematical T operation that inverts the direction of time. Strong and electromagnetic forces are known to be invariant under temporal inversion, however the weak force is not. The *BABAR* experiment recently exploited the quantum-correlated production of B^0 mesons to show that T is a broken symmetry. Here we show that it is possible to perform a wide range of tests of quark flavour changing processes, described by the weak interaction, under the T symmetry in order to validate the Standard Model of particle physics. So while it is unconnected to the entropic arrow of time, the quantum arrow of time sometimes does have a preferred direction, which can be measured.

PACS numbers: 13.25.Hw, 12.15.Hh, 11.30.Er

Weak decays are known to violate the set of discrete symmetries charge conjugation (C), spatial inversion otherwise known as parity (P) [1], CP [2] which distinguishes between matter and antimatter, and time-reversal (T) [3, 4]. These discrete symmetry operations are illustrated in Figure 1. However experimental evidence so far supports the hypothesis that the overall symmetry CPT is conserved, for example see [5–9] for the results of tests using B meson decays. The most significant hint for a non-conservation of CPT comes from *BABAR*, however the CPT violation search using sidereal time evolution of di-lepton decays only has a significance of 2.8σ from expectations of being consistent with the Standard Model of particle physics (SM) [8]. For CPT to be conserved, the level of CP violation has to be balanced by ‘just enough’ T violation so that these two violations cancel each other out to preserve CPT symmetry. While this discussion may seem to be of academic interest it is worth recalling that CPT is conserved in locally invariant field theories such as the SM [10]. However a number of scenarios that have been proposed in order to work toward a theory of quantum gravity can naturally manifest CPT violation. The details of how this might happen depend on the specifics of such models (see for example [11–13]), and as the weak force has a lack of respect for discrete symmetries one should be motivated to test the behaviour of this force with respect to T (and other symmetries) in as many types of weak decay as possible.

The obvious observable to study T symmetry invariance parallels that of a time-dependent or time-integrated CP asymmetry, where the decay of an initial state to a CP eigenstate (or admixture) is compared with the CP conjugate process to test CP . Thus, to test T one can directly compare the probability of the transition of some initial state $|i\rangle$ to some final state $|f\rangle$ and that of the T -conjugate process of the transition of state $|f\rangle$ to




FIG. 1: The effect of T and CP transformations on the set of spatial (x, y) and temporal (t) coordinates. A coordinate system transformed by T results in a temporal inversion, i.e. $t \rightarrow -t$ as shown in the top half of the figure. If the coordinate system is transformed under CP , then matter is changed into anti-matter by C , and spatial coordinates are inverted by P so that $x \rightarrow -x$ and $y \rightarrow -y$. The CPT transformed coordinates result in spatial and temporal inversion.

the state $|i\rangle$, i.e.

$$A_T = \frac{P(|i\rangle \rightarrow |f\rangle) - P(|f\rangle \rightarrow |i\rangle)}{P(|i\rangle \rightarrow |f\rangle) + P(|f\rangle \rightarrow |i\rangle)}. \quad (1)$$

T symmetry invariance is violated if $A_T \neq 0$. The issue that remains is to identify the sets of interesting initial and final states, that can be used for such a test. The decay $(|i\rangle \rightarrow |f\rangle)$ requires a T conjugate partner $(|f\rangle \rightarrow$

$|i\rangle\rangle$) that is experimentally distinguishable.

Historically there have been a number of claims of testing T violation through various methods, and often these tests are not *stand-alone*, but invoke CPT itself to infer that a CP violating asymmetry is telling us about T violation. Such approaches are not able to test T symmetry invariance as they are not able to satisfy the requirement imposed by Eq. (1). In this paper we outline a programme of symmetry invariance tests in B and D decays to extend the remit of the analysis proposed in Ref. [14–16]. We also point out how these measurements relate to the SM weak force quark mixing mechanism given by Cabibbo-Kobayashi-Maskawa (CKM) matrix in the SM [17, 18].

The key to identifying pairs of measurable T conjugate states is highlighted in Ref. [16]. One studies the ensemble of Einstein-Podolsky-Rosen (EPR) [19] correlated neutral meson pairs, and reconstructs an experimentally distinguishable double tag T conjugate pair of transitions: e.g. the flavour of one of the B decays is determined via decay to a flavour specific final state, for example a semi-leptonic B decay, and the other B is a CP tag. Bernabeu et al. proposed the comparison of four sets of processes which satisfy the above criteria:

$$\begin{aligned} \bar{B}^0 &\rightarrow B_- (\ell^+ X, J/\psi K_s^0) \quad \text{vs} \quad B_- \rightarrow \bar{B}^0 (J/\psi K_L^0, \ell^- X) \\ B_+ &\rightarrow B^0 (J/\psi K_s^0, \ell^+ X) \quad \text{vs} \quad B^0 \rightarrow B_+ (\ell^- X, J/\psi K_L^0) \\ \bar{B}^0 &\rightarrow B_+ (\ell^+ X, J/\psi K_L^0) \quad \text{vs} \quad B_+ \rightarrow \bar{B}^0 (J/\psi K_s^0, \ell^- X) \\ B_- &\rightarrow B^0 (J/\psi K_L^0, \ell^+ X) \quad \text{vs} \quad B^0 \rightarrow B_- (\ell^- X, J/\psi K_s^0) \end{aligned}$$

where the K_s^0 and K_L^0 in the final state provide experimentally distinguishable (CP odd and even) T conjugate modes for the CP tag side of the event, and the charge of the lepton identifies the T conjugate flavour tag side.

Recently *BABAR* observed a striking T violation effect via the study of neutral B mesons [4], following the prescription outlined in [16]. The experimental result corresponds to an observation of T violation in these decays at the level of 14σ , which is also consistent with the observed level of CP violation extracted using the same data-set [20]. The Belle experiment should be able to confirm this result, and future flavour factories such as the upgrade of KEKB and Belle II should be able to improve significantly on the precision of this test.

This can be generalised as follows: the ensemble of EPR correlated states produced in e^+e^- interactions at the $\psi(3770)$, $\Upsilon(4S)$, or $\Upsilon(5S)$ is given by

$$\Phi = \frac{1}{\sqrt{2}} \left(P_1^0 \bar{P}_2^0 - \bar{P}_1^0 P_2^0 \right), \quad (2)$$

where $P = B_{d,s}, D$, and the subscript indicates the first or second meson in the pair. At the time one of the mesons decays the wave function collapses into a definite state corresponding to either the first ($P^0 \bar{P}^0$) or the second ($\bar{P}^0 P^0$) ordering. The remaining un-decayed meson will propagate through space-time and mix with its characteristic frequency Δm until it too decays. Given this

one can compute the asymmetry observable of Eq. (1) as a function of the proper time difference Δt between the decay of the first meson t_1 and that of the second meson t_2 . This relies on the quantum coherence of the wave-function of the EPR state over macroscopic scales, which at least in the case of the $\Upsilon(4S)$ has been tested experimentally by Belle [21].

The previous discussion is quite general, the thing that remains is to identify the set of pairs of decay channels that can be used to compute A_T , beyond those already studied. For $B^0(\bar{B}^0)$ meson decays to flavour specific final states, e.g. decays to $X\ell^\pm\nu$, constitute the flavour tag decay part of the problem, and the other decay is into a CP odd or even eigenstate which has an experimentally distinguishable T conjugate final state. It would be possible to use a CP admixture, however one would have the added complication of having to distinguish between CP -even and CP -odd components of an admixture in order to reconstruct orthonormal states of the CP tag decay (for example by performing an angular analysis of neutral pseudo-scalar meson (P) decays to vector-vector final states, and extracting the CP odd part of the transverse amplitude in the decay). For simplicity we mostly restrict further discussion to definite CP final states to illustrate the proposed programme of study.

The measurements performed by *BABAR* used a set of charmonium plus K_s^0 decays ($c\bar{c}K_s^0$) and their T -conjugated partners ($c\bar{c}K_L^0$) paired with flavour tag decays. These $c\bar{c}K^0$ states are theoretically clean to interpret within the SM hence the importance of studying them. The asymmetry parameters measured are $\Delta S_T^\pm$ and $\Delta C_T^\pm$, where

$$A_T \simeq \frac{\Delta C_T^\pm}{2} \cos \Delta m \Delta t + \frac{\Delta S_T^\pm}{2} \sin \Delta m \Delta t. \quad (3)$$

We want to point out that it is possible to perform this measurement with *any* pair of other final states that exhibit time-dependent CP violation such as the T conjugate pairs $B \rightarrow (\eta', \phi, \omega) K_s^0$ and $(\eta', \phi, \omega) K_L^0$. CP tag decays such as the $b \rightarrow s$ loop transitions $\eta' K_s^0$ and $\eta' K_L^0$ are ideal choices for another T violation measurement as physics beyond the SM may affect the asymmetries via non-SM loop amplitudes contributing to the final state. It is important to test the level of T violation in tree processes against that found in loops as a priori the nature of new physics is, by its very definition, unknown. It is possible to study T symmetry invariance in $b \rightarrow u$ transitions via modes such as $B \rightarrow \rho\rho$ and its axial vector counterparts. Here the decay to $\rho^0\rho^0$ is of most interest as there are indications of a sizeable CP odd component in that final state. Similarly one can test $b \rightarrow d$ transitions via $B \rightarrow D^{*+}D^{*-}$ decays. The loop dominated $b \rightarrow s$ decay $B^0 \rightarrow \phi K^*$ is of particular interest as there is a sizeable CP -odd component of the final state, to compare with the dominant T -conjugated CP -even part of the decay. A set of such experimental tests is important as one can independently compare T (and CPT) symmetry violation/invariance for $b \rightarrow u, c, d$, and s transitions

in order to verify if the SM holds up to scrutiny. The time-dependent CP asymmetry parameters have already been determined for these modes, so in a sense half of the job has been done by the B Factories as these experiments have successfully confirmed the real benefit of studying B decays is the miracle that CP violation is manifestly large for these systems.

The asymmetry observables $\Delta S^\pm$ are related to the unitarity triangle angles of the CKM matrix in the SM via $|\Delta S^\pm| = 4\text{Im}\lambda/(1 + |\lambda|^2)$ and $|\Delta C^\pm| = 2(1 - |\lambda|^2)/(1 + |\lambda|^2)$, where the complex parameter λ is related to the mixing parameter ratio q/p and the ratio of amplitudes of the CP tag decay. For the case of the aforementioned $b \rightarrow c, d$, and s quark transitions $\Delta S^\pm$ are measures of $\mp 2\sin 2\beta$, and these are related to the unitarity triangle angle α in the case of $b \rightarrow u$ transitions.

It may be possible to extract information from B_s meson pairs collected at the $\Upsilon(5S)$, however it should be noted that current vertex detector technology is insufficient to resolve individual neutral meson oscillations in this system at an e^+e^- collider, so one must rely on the $\Delta\Gamma$ modulation of the neutral meson oscillation to obtain information related to A_T .

Now we can turn to the issue of charm decays where we recently urged the experimental community to embark upon the systematic study of time-dependent CP asymmetries to parallel the work of the B Factories since 1999 [22]. This was in part motivated by the opportunity to make such measurements given the availability of data from LHCb and the promise of more to come from future flavour factories. Following our previous paper there have been tantalising indications of a non-zero direct CP asymmetry in $D \rightarrow K^+K^-$ and $D \rightarrow \pi^+\pi^-$ decays from LHCb and CDF [23, 24] that increase the urgency of performing these tests. The underlying physics regarding the production of neutral D mesons in e^+e^- collisions at the $\psi(3770)$ is a direct parallel of the production of B mesons at the $\Upsilon(4S)$. We note that as a consequence of this, one can re-use the measurement technique of Ref. [16] adopted by *BABAR* and apply this to T violation searches in charm mesons, with the caveat that the lifetime difference matters for charm as $y = \Delta\Gamma/2\Gamma$ is non-zero. In analogy with our observation that there is a set of CP -eigenstate or admixture modes that should be studied for T symmetry invariance in B decays, at an asymmetric τ -charm factory running at the $\psi(3770)$ one also has a set of interesting decay modes to explore. Thus one can study T invariance for $c \rightarrow d$ and $c \rightarrow s$ transitions at leading order, and also in principle access $c \rightarrow b$ transitions in terms of the second order, CKM suppressed, loop contributions given sufficient data. This last transition is related to observables reported in the $B \rightarrow c\bar{c}K^0$ analysis by *BABAR*.

In the SM, the level of CP violation in charm is expected to be small so, unless T violation arises from other mechanisms, any experimental test of T symmetry invariance would require a high degree of systematic control (and careful design to minimise systematic un-

certainties), and a large integrated luminosity of data collected via $e^+e^- \rightarrow \psi(3770) \rightarrow D^0\bar{D}^0$. Given the sample sizes of experiments under discussion, with a few ab^{-1} , one would not expect to be sensitive to the level of T violation compatible with the SM — however that does mean that if one were to perform a T symmetry measurement and observe T violation, that would have to result from physics beyond the SM. Experimentally, the dilution effect present in determining flavour tag B states is absent for semi-leptonic flavour tagged neutral D mesons as there are only primary leptons produced in $D^0(\bar{D}^0) \rightarrow X_s\ell\nu$ decays, and as both D mesons are reconstructed with a well known initial set of conditions, there should be very little background present (at least in final states without a K_L^0 meson) with which one could study T symmetry invariance. The other experimental issue of relevance is that of detector resolution. Using vertex detector technology accessible today, with a center of mass boost factor $\beta\gamma$ relative to the laboratory frame of at least about 0.3-0.4, it should be possible for a τ -charm factory to study T symmetry invariance in a wide range of final states including $h^+h^-K_{s,L}^0$. CP tag decays such as $D^0 \rightarrow K_{s,L}^0\pi^0$ should be experimentally accessible in terms of CP violation measurements, however the lack of charged particles originating from the D decay vertex means that the tag mode with $CP = +1$ would have to be reconstructed via π^0 Dalitz decays or photon conversion processes. However equivalent measurements with $D^0 \rightarrow K_{s,L}^0(\omega, \eta, \eta', \rho^0, \phi, f_0, a_0)$ are also possible, where these are expected to be less experimentally challenging than the $K_{s,L}^0\pi^0$ combination. Studies of $D^0 \rightarrow K_s^0K_s^0K_L^0$ and the T -conjugate $K_L^0K_L^0K_s^0$ mode could be used to explore these effects for W exchange amplitudes, if one could suitably reconstruct the decay vertex of the latter mode. These allow one to probe T symmetry invariance in $c \rightarrow d$ and $c \rightarrow s$ transitions to complement the set of measurements in b quark transitions.

In summary, the recent observation of T violation in $b \rightarrow c$ transitions by *BABAR* raises an interesting (and old) question: what discrete symmetries are respected by the weak force? We propose a number of measurements that can be made to test T symmetry invariance in $b \rightarrow u, d$ and s transitions (to complement the existing $b \rightarrow c$ tests), as well as instigating a programme of measurements in the charm sector to test $c \rightarrow d$ and $c \rightarrow s$ transitions. One expects sizeable T violation in the various B decay final states to balance the open form of the bd Unitarity triangle, and small effects (essentially zero within experimental precision that would be achievable with facilities under consideration today) for D decays. Any observed T violation can be related directly to the SM via existing knowledge of the CKM matrix, and so one can systematically probe the behaviour of the fundamental weak force in terms of the T symmetry to complement almost five decades of study with regard to CP . The $b \rightarrow d$ and s transition effects should be comparable in magnitude to those reported by *BABAR* for $b \rightarrow c$

transitions as they are a measure of the unitarity triangle angle β , whereas one expects a smaller value for $b \rightarrow u$ transitions which are a measure of α in the SM. By measuring this set of decays one would be able to constrain leading and higher order T violation contributions in the SM to complement the set of CP violation constraints reported by the B Factories since 1999. To do this one requires high statistics data samples from future e^+e^- based (asymmetric energy) B and τ -charm flavour factories operating at the $\Upsilon(4S)$ in the case of b quark transitions and the $\psi(3770)$ in the case of c quark transitions. The data samples recorded by *BABAR* and Belle provide a starting point for detailed exploration of T violation in B decays before Belle II starts taking data later this decade. The full set of measurements indicated in this

article would probe T violation in tree and loop transitions for both up and down type quarks. It is also worth remembering that one can probe CPT in an analogous way to over-constrain the model of quark-flavour changing weak interactions, namely the Kobayashi-Maskawa mechanism and the CKM matrix.

I. ACKNOWLEDGMENTS

We wish to thank Jose Bernabeu, Fernando Martinez-Vidal and Brian Meadows for useful discussions during the preparation of this paper.

-
- [1] C. S. Wu, E. Ambler, R. W. Hayward, D. D. Hoppes and R. P. Hudson, Phys. Rev. **105**, 1413 (1957).
 - [2] J. H. Christenson, J. W. Cronin, V. L. Fitch and R. Turlay, Phys. Rev. Lett. **13**, 138 (1964).
 - [3] A. Angelopoulos *et al.* [CLEAR Collaboration], Phys. Lett. B **444**, 43-51 (1998).
 - [4] J. P. Lees *et al.* [BABAR Collaboration], Phys. Rev. Lett. **109**, 211801 (2012) [arXiv:1207.5832 [hep-ex]].
 - [5] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. Lett. **92**, 181801 (2004) [hep-ex/0311037].
 - [6] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. D **70**, 012007 (2004) [hep-ex/0403002].
 - [7] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. Lett. **96**, 251802 (2006) [hep-ex/0603053].
 - [8] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. Lett. **100**, 131802 (2008) [arXiv:0711.2713 [hep-ex]].
 - [9] T. Higuchi *et al.* [Belle Collaboration], Phys. Rev. D **85**, 071105 (2012) [arXiv:1203.0930 [hep-ex]].
 - [10] G. Luders, Kong. Dan. Vid. Sel. Mat. Fys. Med. **28N5**, 1 (1954).
 - [11] V. A. Kostelecky, hep-ph/0202094.
 - [12] J. Bernabeu, N. E. Mavromatos and J. Papavassiliou, Phys. Rev. Lett. **92**, 131601 (2004) [hep-ph/0310180].
 - [13] J. Bernabeu, N. E. Mavromatos and S. Sarkar, Phys. Rev. D **74**, 045014 (2006) [hep-th/0606137].
 - [14] M. C. Banuls and J. Bernabeu, Phys. Lett. B **464**, 117 (1999) [hep-ph/9908353].
 - [15] M. C. Banuls and J. Bernabeu, Nucl. Phys. B **590**, 19 (2000) [hep-ph/0005323].
 - [16] J. Bernabeu, F. Martinez-Vidal and P. Villanueva-Perez, JHEP **1208**, 064 (2012) [arXiv:1203.0171 [hep-ph]].
 - [17] N. Cabibbo, Phys. Rev. Lett. **10**, 531-533 (1963).
 - [18] M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652-657 (1972).
 - [19] A. Einstein, B. Podolsky, N. Rosen Phys. Rev. **47**, 777-780 (1935).
 - [20] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. D **79**, 072009 (2009) [arXiv:0902.1708 [hep-ex]].
 - [21] A. Go *et al.* [Belle Collaboration], Phys. Rev. Lett. **99**, 131802 (2007) [quant-ph/0702267 [QUANT-PH]].
 - [22] A. J. Bevan, G. Inguglia and B. Meadows, Phys. Rev. D **84**, 114009 (2011) [arXiv:1106.5075 [hep-ph]].
 - [23] R. Aaij *et al.* [LHCb Collaboration], Phys. Rev. Lett. **108**, 111602 (2012) [arXiv:1112.0938 [hep-ex]].
 - [24] T. Aaltonen *et al.* [CDF Collaboration], Phys. Rev. Lett. **109**, 111801 (2012) [arXiv:1207.2158 [hep-ex]].